

COLONOSCOPY [SUTAB Pill Prep]

This is a split-dose (2-day) regimen. A total of 24 tablets is required for complete preparation for colonoscopy. You will take the tablets in two doses of 12 tablets each. Water must be consumed with each dose of SUTAB, and additional water must be consumed after each dose.

ON THE DAY BEFORE YOUR PROCEDURE

You **MAY HAVE** a low residue breakfast. Low residue foods include eggs, white bread, cottage cheese, yogurt, grits, coffee, and tea. You may have clear liquids.

You **CANNOT HAVE**

- milk or eat or drink anything colored red or purple.
- alcohol.
- other laxatives while taking SUTAB.
- oral medications within 1 hour of starting each dose of SUTAB.
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, or penicillamine, take these medications at least 2 hours before and not less than 6 hours after administration of each dose of SUTAB.

Liquids That Are OK to Drink

- Coffee or tea (no cream or nondairy creamer)
- Fruit juices (without pulp)
- Gelatin desserts (no fruit or topping)
- Water
- Chicken broth
- Clear soda (such as ginger ale)

ON THE DAY PRIOR TO COLONOSCOPY [Dose 1]

STEP 1 Open 1 bottle of 12 tablets.

STEP 2 Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water and drink the entire amount of water over 15 to 20 minutes. **NOTE:** *You do not have to swallow all 12 pills at the same time. Take 1 or 2 pills every couple of minutes until they are gone.*

IMPORTANT: If you experience preparation-related symptoms (for example, nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until your symptoms diminish.

Drink additional water

STEP 3 Approximately 1 hour after the last tablet is ingested, fill the provided container again with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.

STEP 4 Approximately 30 minutes after finishing the second container of water, fill the provided container with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.

DAY OF THE COLONOSCOPY [Dose 2]

- Continue to consume only clear liquids until after the colonoscopy.
- The morning of the colonoscopy (5 to 8 hours prior to the colonoscopy and no sooner than 4 hours from starting Dose 1), open the second bottle of 12 tablets.
- Repeat **STEP 1** to **STEP 4** from **DOSE 1**.

